

Title:

Challenging the gold standard: endohepatology for the diagnosis of presinusoidal portal hypertension

Authors:

Sofia Bragança, Pooja Khonde, Stephen D. Zucker's, Marvin Ryou

DOI: 10.17235/reed.2025.11480/2025

Link: [PubMed \(Epub ahead of print\)](#)

Please cite this article as:

Bragança Sofia, Khonde Pooja, Zucker's Stephen D., Ryou Marvin. Challenging the gold standard: endohepatology for the diagnosis of presinusoidal portal hypertension. Rev Esp Enferm Dig 2025. doi: 10.17235/reed.2025.11480/2025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CC 11480

Challenging the gold standard: endohepatology for the diagnosis of presinusoidal portal hypertension

Sofia Bragança¹, Pooja Khonde^{2,4}, Stephen D. Zucker's^{3,4}, Marvin Ryou^{3,4}

¹Department of Gastroenterology. Hospital Professor Doutor Fernando Fonseca EPE. Amadora, Portugal. Departments of ²Pathology and ³Gastroenterology, Hepatology and Endoscopy. Brigham and Women's Hospital. Boston, Massachusetts. United States. ⁴Harvard Medical School. Boston, Massachusetts. United States

Correspondence: Sofia Bragança

e-mail: sofia_58632@hotmail.com

Authors' contribution: Supervision: M. R.; visualization: P. K., S. D. Z.; writing-original draft: S. B.

Financial support: The corresponding author received financial support through the 2024 Digestive Endoscopy Internship Grant from the Núcleo de Gastreenterologia dos Hospitais Distritais (NGHD).

Informed consent: The authors obtained informed consent from the patient for the publication of their information and imaging.

Conflict of interest: Marvin Ryou, medical consultant for Olympus, Fujifilm and Cook Medical.

Artificial intelligence: The authors declare that they did not use artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article.

Keywords: Endoscopic ultrasound. Portal pressure gradient. Liver biopsy. Shear wave elastography. Nodular regenerative hyperplasia.

Dear Editor,

We report the case of a 52-year-old patient referred for evaluation of abnormal liver tests, with extensive etiological workup that was negative. Abdominal computed tomography (CT) and magnetic resonance imaging (MRI) revealed hypertrophy of the left hepatic lobe, nodular liver contours, and ascites (serum ascites albumin gradient [SAAG] 1.7; total protein 1.8 g/dl). Upper endoscopy showed no stigmata of portal hypertension.

A dual interventional radiology procedure was performed. Transjugular liver biopsy (LB) (18 G) was non-diagnostic, and hepatic venous pressure gradient (HVPG) was normal (HVPG 3 mmHg; wedged hepatic venous pressure [WHVP] 13 mmHg; free hepatic venous pressure [FHRVP] 10 mmHg).

Given the suspicion of pre-sinusoidal portal hypertension, endoscopic ultrasound (EUS) was performed (Arietta™ 850, Fujifilm Healthcare, Tokyo, Japan) (Fig. 1A-F). EUS showed blunting of the left lobe tip, irregular contours, and ascites. Endoscopic “palpation” revealed an indentation of 2.0 mm (abnormal). Shear-wave-elastography (SWE) of the left lobe was normal at 6.04 kPa, but other areas showed 10.5-11.1 kPa (abnormal). There was no splenomegaly, but the mean SWE was 31.9 kPa (abnormal).

EUS-guided portal pressure gradient (EUS-PPG), measured with EchoTip Insight™ FNB needle (Cook Medical, USA), was elevated (PPG 8 mmHg; mean hepatic vein pressure 4.0 mmHg; mean portal vein pressure [PVP] 12.0 mmHg). Targeted EUS-guided biopsy of the left lobe (Acquire™ 19G FNB needle, Boston Scientific, USA) showed vague liver nodular transformation with alternating regenerative and atrophic hepatocytes, consistent with nodular regenerative hyperplasia (NRH) (Fig. 2A and B).

Due to refractory ascites, transjugular intrahepatic portosystemic shunt (TIPS) placement was proposed. Pre-TIPS evaluation showed mildly elevated HVPG (6 mmHg), while immediate post-TIPS assessment revealed elevated HVPG (HVPG 11 mmHg; PVP 20 mmHg; FHRVP 9 mmHg). Ascites was controlled with TIPS.

Discussion

Endohepatology has emerged in recent years as a means to enhance the interface between endoscopy and hepatology, allowing a “one-stop-shop” evaluation in selective patients. Nowadays, several applications have been integrated into clinical practice of some centers (1,2). Among them, two stand out for their relevance, the measurement of the PPG and LB, the latter allowing single-operated targeted-LB (3,4). Additionally, detailed assessment of the hepatic parenchyma, indentation and hepatic and splenic SWE, enhance its value as a diagnostic tool. Together, these techniques have significantly advanced the evaluation of portal hypertension, particularly in cases where conventional approaches may be inconclusive (5).

There is no head-to-head studies currently available to definitely establish the superiority of PPG over HVPG. Nevertheless, several advantages have been reported, particularly in the context of pre-sinusoidal portal hypertension, as in NRH, because HVPG may underestimated the portal pressure (4).

References

1. Diehl DL. The four pillars of endohepatology. J Can Assoc Gastroenterol 2025;8:56-61. DOI: 10.1093/jcag/gwae036
2. Bazarbashi AN, Al-Obaid L, Ryou M. Future directions in endohepatology. TIGE 2022;24:98-107. DOI: 10.1016/j.tige.2021.09.006
3. Diehl DL. Top tips regarding EUS-guided liver biopsy. Gastrointest Endosc 2022;95:368-71. DOI: 10.1016/j.gie.2021.10.016
4. Bazarbashi AN, Ryou M. Portal pressure measurement: Have we come full circle? Gastrointest Endosc 2021;93:573-6.
5. Choi AY, Chang KJ. Endoscopic diagnosis of portal hypertension. TIGE 2022;24:167-75. DOI: 10.1016/j.tige.2022.01.003




Fig. 1. One-stop-shop endoscopic ultrasound (EUS) approach. A. Endoscopic ultrasound (EUS) liver observation suggestive of advanced chronic liver disease. B. Abnormal EUS-liver indentation. C. Normal left lobe shear-wave elastography. D. Abnormal left lobe shear-wave-elastography. E. EUS-guided portal pressure gradient (PPG) - portal vein. F. PPG measurement - left hepatic vein.




Fig. 2. Histopathological examination. Hematoxylin and eosin (A) and reticulin stain (B), 20x: nodular liver parenchyma secondary to atrophic hepatocytes (arrows) flanking regenerating, hypertrophic hepatocytes. There is no fibrosis.