

**Title:**

**Magnesium poisoning, multi-organ failure and ischemic colitis secondary to chronic ingestion of a parapharmacy product**

**Authors:**

Carmen Díaz Gómez, Lucía López Amor, Emilio García Prieto, Dolores Escudero Augusto

DOI: 10.17235/reed.2018.5355/2017

Link: [PubMed \(Epub ahead of print\)](#)

Please cite this article as:

Díaz Gómez Carmen, López Amor Lucía, García Prieto Emilio, Escudero Augusto Dolores. Magnesium poisoning, multi-organ failure and ischemic colitis secondary to chronic ingestion of a parapharmacy product. Rev Esp Enferm Dig 2018. doi: 10.17235/reed.2018.5355/2017.

Enero 2017 • Volumen 109 • Número 1 • Páginas 1-86

CODE: REED09 ISSN: 1000-0188

Revista Española de Enfermedades Digestivas  
THE SPANISH JOURNAL OF GASTROENTEROLOGY

Acceso al texto completo en: [www.reed.es](http://www.reed.es) y [www.sped.es](http://www.sped.es)

Factor de Impacto 100 (IIF): JCR: 1.455 (2016)  
SCR: 0.34 (2016)

ORGANO OFICIAL DE:  
SOCIEDAD ESPAÑOLA DE PATOLOGÍA DIGESTIVA, SOCIEDAD ESPAÑOLA DE ENDOSCOPIA DIGESTIVA Y ASOCIACIÓN ESPAÑOLA DE ECOGRAFÍA DIGESTIVA

SEPO

<b>Editorial</b> Colitis diverticular bleeding. Have we identified the risk factors for massive bleeding yet? J. W. Barcia Valencia	<b>Lesión de Bowdler: diagnóstico por gastroscopia</b> R. Barrio Guea, N. N. Barrio Irujo, M. Paz Noya y J. E. Domínguez Muñoz	65
<b>Trabajos Originales</b> Risk factors for severity and recurrence of colonic diverticular bleeding N. Argente, P. Cabero, A. Arduas, M. Escobar and N. Guzmán	<b>Notas Clínicas</b> Celiac crisis in adults: a case report and review of the literature focusing in the prevention of relapsing syndrome M. de Alaveda-Nemón, V. L. Barrio-Cabral and S. L. Linares	67
Management and inflammatory bowel disease: the other diagnosis J. Barrio, A. Argente, V. L. Barrio-Cabral, J. Castro, M. Acari-Barris, M. Hernández-Serra, C. García, M. de Alaveda-Nemón and A. López-García	Herangopercistoma patético. Una localización infrecuente de tumor estomacal I. Alamo Abad, J. M. García-Cerbal, L. Aguirre-Duñabe, A. M. Quintana Berro y A. Calvo Artero	69
Influence of sustained 48-hr response on the regression of fibrosis and portal hypertension in cirrhotic HCV patients treated with antiviral therapy A. Barrio, J. Cabero, M. J. López-Alcalá, I. Fariña, M. T. Arce, A. Galván, F. Castiella, E. Fabrega and J. Crespo	Hemólisis por eritropoiesis suprarrenal C. Pizarro-Carral, A. Escobedo-Sánchez, M. A. Paredes-Capó, J. Arangul-Abad y C. García-Delgado	70
Malnutrition risk questionnaire combined with body composition measurement in malnutrition screening in inflammatory bowel disease A. A. Cortés, A. Muñoz, Z. Pili, I. Pall and P. Muñoz	Endoscópico removal of intubated large sigmoid polyp: a case report M. Escudero Augusto and S. López-Munoz	73
A survey-based analysis on endoscopic quality indicators compliance among Spanish endoscopists I. Fernández-Cruz, F. Argente, P. Alonso, J. Soto and S. Soriano	Meconíosis obstruente en neonato: caso de abdomen agudo A. Tapia-Palacio, M. R. Ramos-Vázquez, J. C. Cordero-Ramos, J. Cordero-Lafont and D. L. Carballo-Pérez	76
<b>Revisión</b> Endoscopic resection of colonic polyps in patients on antiplatelet therapy: an evidence-based guideline for clinicians G. Piana, M. Sostero-Salín, C. Salinas, F. Day and M. J. Cuatrecasas	<b>Cartas al Editor</b> Neoplasia neuroendocrina intestinal, un tumor poco habitual M. de Barrio Noya, J. Barrio Fernández y M. N. Barrio Rodríguez	79
<b>Indicadores en Patología Digestiva</b> Trastorno de la arteria mesentérica superior: una causa infrecuente de obstrucción intestinal J. Sempere Jaque, P. Alamo Barrio y J. C. García-Pérez	Presentación intestinal de tuberculosis por radioterapia pélvica: diagnóstico asociado a cambios de Baxant, patología infrecuente y poco conocida Y. Pina-Vargas, D. M. Aguirre y L. A. Alvarado	80
Neumatoxina sintética intravital A. F. Romero-Muñoz y K. Barrio-Zelga	Parrot endoscopy assistance for an achalasia patient with multiple esophageal diverticula Y. Se, H. Zhu and D. Liu	81
Trastorno de Hirsch a cinco años de edad C. Oña-Solís, C. C. Hernández-Segura, P. Pineda-Rodríguez y A. N. González-Fernández	Actualización y endoscopia: endoscopia infundibular en el diagnóstico diferencial de abdomen agudo C. Sánchez-Jiménez, J. Guevara-Nieto y J. A. Acosta-Paredes	81
Endoscópico removal of leiomyomas in a scleroderma patient J. L. Barrio-Hernández, M. E. Torres-Castro and M. Torres-Rodríguez	Perforación múltiple de divertículos de intestino delgado en paciente con síndrome de Down R. Hernández-Cabrera, A. Barrio-Cabrera y E. Palencia-López	83
All that glitters is not gold. A different cause for an "alkaline colitis" A. Pineda, M. Soto, J. Villaverde and G. Navarro	Altoplexia analítica como manifestación paraneoplásica de un adenoma actínico gástrico J. Barrio-Cabrera, F. Fernández-Serrano y J. de la Fuente-Aguado	83
	<b>Revisores 2016</b>	85

ARAN

[www.reed.es](http://www.reed.es)

*This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.*

**NC 5355 inglés**

**Magnesium poisoning, multi-organ failure and ischemic colitis secondary to chronic ingestion of a parapharmacy product**

Carmen Díaz-Gómez<sup>1</sup>, Lucía López-Amor<sup>1</sup>, Lorena Forcelledo-Espina<sup>1,2</sup>, Emilio García-Prieto<sup>1,3,4</sup> and Dolores Escudero-Augusto<sup>1,2</sup>

<sup>1</sup>Intensive Medicine Service. Hospital Universitario Central de Asturias. Oviedo, Asturias. Spain. <sup>2</sup>Critical Patient Translational Research Group. Instituto de Investigación Sanitaria del Principado de Asturias (IISPA). Oviedo, Asturias. Spain. <sup>3</sup>Department of Medicine. University of Oviedo. Oviedo, Asturias. Spain. <sup>4</sup>Translational Microbiology Group. Instituto de Investigación Sanitaria del Principado de Asturias (IISPA). Oviedo, Asturias. Spain

**Received:** 06/11/2017

**Accepted:** 06/02/2018

**Correspondence:** Carmen Díaz Gómez. Intensive Medicine Service. Hospital Universitario Central de Asturias. Av. de Roma, s/n. 33011 Oviedo, Asturias  
e-mail: carmendiazg@icloud.com

**ABSTRACT**

The article describes the clinical case of a patient with multi-organ failure and ischemic colitis secondary to magnesium poisoning, due to the chronic intake of a parapharmacy product, used for the treatment of chronic constipation. The clinical case is described and a review of the literature is made.

**Key words:** Magnesium. Iatrogenic disease. Renal replacement therapy. Enterocolitis. Ileitis.

**INTRODUCTION**

Hypermagnesemia is defined as a serum magnesium level greater than 1 mmol/l (1). The cause is usually iatrogenic and there is a higher risk associated with elderly

patients, chronic renal failure with reduced excretion and gastrointestinal disorders with increased absorption (2,3).

Mild hypermagnesemia can be asymptomatic (4) and symptoms start to appear when the levels are  $> 2.5$  mmol/l (1), as shown in table 1. A relationship has been described between hypermagnesemia and the need for vasopressor drugs, a higher risk of respiratory failure and an increased mortality (5-7).

Hypermagnesemia treatment depends on the levels and is based on stopping magnesium administration, increasing the excretion by loop or thiazide diuretics, proper fluid administration and calcium salts to protect the myocardium. Severe cases require renal replacement to decrease magnesium levels to a non-toxic level in a short period of time (1).

## **CASE REPORT**

The case was a 70-year-old woman with a history of mantle cell lymphoma, pudendal neuralgia and painful bladder syndrome treated with opioid and intravesicular botulinum toxin. The last dose was administered ten months prior to being admitted. Furthermore, she had a history of chronic constipation, even before starting opioid treatment, and was taking magnesium in a granulated formulation from a parapharmacy. The product was Magnesium Carbonate<sup>®</sup>, which contained 290 mg of magnesium per each gram, and the patient was taking one tablespoon per day. The product label recommended to “dissolve two tablespoons in half a glass of water or juice after meal digestion and before bedtime”. There was no other information or cautions in the case of an overdose.

The patient developed vomiting and weakness and was initially treated at home by the extra hospital Emergency Service. She was alert and orientated (Glasgow coma scale of 14 point), with a blood pressure (BP) of 104/57 mmHg and paresis in the four limbs with a muscle strength of 1/5. She also presented areflexia, respiratory failure with tachypnea of 32 breaths/min and an oxygen saturation of 83%. Non-invasive mechanical ventilation was applied with an improvement in oxygen saturation to 98%. During the medical interview, the patient admitted to taking magnesium carbonate as a laxative on a daily basis. Therefore, magnesium intoxication was suspected and treatment with fluids was started at 0.5 mg IV of calcium chloride and 20 mg of

furosemide. The patient was transferred to the Emergency Room of her reference hospital and remained hemodynamically unstable (BP 98/64 mmHg) despite fluid administration, and was therefore treated with vasoactive drugs.

The muscular weakness turned into an areflexic tetraplegia that required non-invasive mechanical ventilation support to maintain oxygen saturation and the blood gas analysis within normal values. Serum magnesium levels were 7.35 mmol/l and therefore energetic fluid therapy plus 1 g of calcium gluconate was administered. The patient was transferred to the Intensive Care Unit (ICU) and became more hemodynamically unstable (BP 74/37 mmHg) with oligoanuric renal failure and respiratory failure due to the tetraplegia, requiring intubation and invasive mechanical ventilation. Given the severity of the case, hemodiafiltration was performed. Despite a decrease in the serum magnesium level to 4.55 mmol/l after three hours of renal replacement, the shock persisted and noradrenaline at  $> 1.5$  mcg/kg/min was required. The patient had severe metabolic acidosis and lactate levels of 9.7 mmol/l. An echocardiography was performed and there were no relevant findings due to a poor echographic window.

During the first few hours after admission, the patient developed an important abdominal distension with massive diarrhea. An abdominal-pelvic computed tomography (CT) scan identified a severe dilatation affecting the entire colon without signs of intestinal ischemia (Fig. 1). These findings were considered as a toxic megacolon secondary to a magnesium intoxication.

After ten hours of renal replacement, the serum magnesium level dropped to 3.1 mmol/l. The hemodynamic situation progressively improved and the magnesium levels decreased in parallel. The noradrenaline infusion was stopped 72 h after admission. However, abdominal distension and diarrhea persisted, accompanied by a fever of 38.7 °C. New complications were discarded after an abdominal CT scan on the 4<sup>th</sup> day of the clinical course. Afterwards, clinical improvement continued and extubation was achieved on the 7<sup>th</sup> day. The patient was discharged from the ICU on the 10<sup>th</sup> day of the clinical course.

A colonoscopy was performed due to persistent diarrhea that identified signs of ischemic colitis on a 30 cm segment length of the colon (Fig. 2). The abdominal

affectation recovered very slowly and hospital discharge was delayed until the 67<sup>th</sup> day.

## **DISCUSSION**

Magnesium salts are commonly used as osmotic laxatives for patients suffering from chronic constipation (8) and is recommended by both gastroenterology specialists and general practitioners due to its proven efficacy (9). However, they are not always prescribed by health professionals as they are accessible in parapharmacies. A chronic intake of magnesium products with no medical control can lead to severe intoxications that can cause multiple organ failure and death (Table 1) especially in people with the risk factors mentioned previously such as elderly cases, chronic renal failure and gastrointestinal disorders.

In the most severe gastrointestinal manifestations, hypermagnesemia can cause toxic megacolon and ischemic colitis due to an altered intestinal blood flow. This leads to a decrease in bowel motility caused by fecal impaction and an increase in the intraluminal pressure exacerbated by a prolonged hypotension (10), as in this case.

Parapharmacy products are not considered as “drugs” by the consumer. Thus, patients have a false feeling that they can safely take this medication in an uncontrolled manner and do not mention this during the medical interview. Furthermore, the posology information of these products can be ambiguous and does not alert the consumers to the risk of overdose.

This case report highlights the risks of self-medication with parapharmacy products that contain magnesium. Therefore, when intoxication is suspected, this etiology should be considered and assessed via a rigorous medical interview.

## **REFERENCES**

1. Torregrosa JV, Torres A. Alteraciones del metabolismo del calcio, del fósforo y del magnesio. En: Farreras Rozman. Medicina Interna. XVII Ed. Elsevier España; 2016. pp. 799-807.
2. Kala J, Abudayyeh A. Magnesium: an overlooked electrolyte. J Emerg Med 2017;52(5):741-3. DOI: 10.1016/j.jemermed.2017.01.011

3. Karahan MA, Kuckuc A, Buyukfirat E, et al. Acute respiratory and renal failure due to hypermagnesemia, induced by counter laxatives in an elderly man. *J Clin Diagn Res* 2015;9(12):UL01. DOI: 10.7860/JCDR/2015/16630.6939
4. Yoon HE, Kim YW, Ha KS, et al. Hypermagnesemia accompanied with colonic perforation in a hemodialysis patient. *Yonsei Med J* 2013;54:797-800. DOI: 10.3349/ymj.2013.54.3.797
5. Onish S, Yoshino S. Cathartic-induced fatal hypermagnesemia in the elderly. *Intern Med* 2006;45:207-10. DOI: 10.2169/internalmedicine.45.1482
6. Celi LA, Scott DJ, Lee J, et al. Association of hypermagnesemia and blood pressure in the critically ill. *J Hypertens* 2013;31:2136-41. DOI: 10.1097/HJH.0b013e3283642f18
7. Thongprayoon C, Cheungpasitporn W, Srivli N, et al. Admission serum magnesium levels and the risk of acute respiratory failure. *Int J Clin Pract* 2015;69:1030-8. DOI: 10.1111/ijcp.12696
8. Cheungpasitporn W, Thongprayoon C, Quian Q. Dymagnesemia in hospitalized patients: prevalence and prognostic importance. *Mayo Clin Proc* 2015;90:1001-10. DOI: 10.1016/j.mayocp.2015.04.023
9. Weng YM, Chen SY, Chen HC, et al. Hypermagnesemia in a constipated female. *J Emerg Med* 2013;44:57-60. DOI: 10.1016/j.jemermed.2011.09.004
10. Sugiyama M, Kusumoto E, Ota M, et al. Induction of potentially lethal hypermagnesemia, ischemic colitis, and toxic megacolon by a preoperative mechanical bowel preparation: report of a case. *Surg Case Rep* 2016;2(1):18. DOI: 10.1186/s40792-016-0145-6

**Table 1. Symptoms according to magnesium levels**

<i>Serum magnesium levels</i>	<i>Symptoms</i>
2-3 mmol/l	Nausea, vomiting, facial flushing, headache, hyporeflexia, lethargy
3-5 mmol/l	Drowsiness, hypocalcemia, loss of deep tendon reflexes, hypotension, bradycardia, changes in the ECG
> 5 mmol/l	Muscular/muscle paralysis, respiratory depression, complete heart blockage, coma, cardiac arrest

ECG: electrocardiography.




Fig. 1. Abdomino-pelvic CT with contrast in the coronal (A) and axial (B) sections showing a severely dilated colon, fecal and fluid retention without signs of obstruction, pneumatosis or an intestinal loop.





Fig. 2. Colonoscopy showing edema, erythema and ulcers covered by fibrin related to ischemic colitis. The changes were mostly circumferential and more severe towards the distal area.

Accepted Article