

Title:
Type V biliary cyst with cystolithiasis

Authors:
Martín Bailón Cuadrado, Pilar Pinto
Fuentes, David Pacheco Sánchez, Trinidad
Escudero Caro

DOI: 10.17235/reed.2018.5477/2018

Link: [PubMed \(Epub ahead of print\)](#)

Please cite this article as:
Bailón Cuadrado Martín, Pinto Fuentes
Pilar, Pacheco Sánchez David, Escudero
Caro Trinidad. Type V biliary cyst with
cystolithiasis. Rev Esp Enferm Dig 2018.
doi: 10.17235/reed.2018.5477/2018.

Enero 2017 • Volumen 109 • Número 1 • Páginas 1-86

CODE: REED09N ISSN: 1009-0108

Revista Española de Enfermedades Digestivas
THE SPANISH JOURNAL OF GASTROENTEROLOGY

Acceso al texto completo en: www.reed.es o www.sped.es

Factor de Impacto 100 (I-I): 1.455 (2016)
SCR: 0.34 (2016)

ORGANO OFICIAL DE:
SOCIEDAD ESPAÑOLA DE PATOLOGÍA DIGESTIVA, SOCIEDAD ESPAÑOLA
DE ENDOSCOPIA DIGESTIVA Y ASOCIACIÓN ESPAÑOLA DE ECOGRAFÍA DIGESTIVA

SEPO

Editorial Celiac disease: Have we identified the risk factors for massive bleeding yet? J. W. Berca Valencia	Lesión de Bouveret: diagnóstico por gastroscopia R. Barrio Guea, N. N. Barrio Irujo, M. Paz Noya y J. E. Dominguez Muñoz	65
Trabajos Originales Risk factors for severity and recurrence of colonic diverticular bleeding N. Azupard, P. Cabero, A. Arduini, M. Escobar and N. Guzmán	Notas Clínicas Celiac crisis in adults: a case report and review of the literature focusing on the presentation of subacute syndrome M. de Alaveda-Nemón, V. L. Barrio-Caballero and S. Latorre	67
Microsporidian and inflammatory bowel disease: the other diagnosis J. Bermejo, A. Rojas, S. López-Cerdá, J. Cuervo, M. Acuña-Barca, M. Hernández-Serna, C. García, M. de la Cruz, D. Barrio and A. López-Soriano	Herangopercistoma patiens. Una localización infrecuente de tumor intestinal I. Alonso Abad, J. M. García-Cerco, L. Aguirre-Díaz, A. M. Quintana Berro y A. Colla-Morales	69
Influence of sustained blood response on the regression of fibrosis and portal hypertension in cirrhotic (HCV) patients treated with antiviral therapy A. Barrio, J. Cabero, W. J. López-Alcalá, I. López, M. T. Ariza, A. Galván, F. Castiella, E. Fabrega and J. Crespo	Hemostasia por erupción papilar intraductal C. Pizarro-Carpas, A. Escobedo-Sánchez, M. A. Paredes-Capó, J. Arangul-Arribas y C. García-Delgado	70
Malnutrition risk questionnaire combined with body composition measurement in malnutrition screening in inflammatory bowel disease A. A. Cortés, A. Muñoz, Z. Pili, I. Pall and P. Muñoz	Endoscopic removal of intubated large varicose gastric: a case report M. Oquendo and J. Sainza-Munizaga	73
A survey-based analysis on endoscopic quality indicators compliance among Spanish endoscopists I. Fernández-Cruz, F. Argüelles, P. Alonso, J. Salas and S. Soriano	Meconitosis obstruente en neonato: caso de abdomen agudo A. Tapia-Palacio, M. R. Ramos-Vázquez, J. C. Cordero-Ramos, J. Cordero-Lafont and L. Carballo-Pérez	76
Revisión Endoscopic resection of colorectal polyps in patients on antiplatelet therapy: an evidence-based guideline for clinicians G. Piana, M. Sostero-Salín, C. Salinas, F. Day and M. J. Cuatrecasas	Cartas al Editor Neoplasia neuroendocrina intestinal, un tumor poco habitual M. de Barrio Noya, J. Barrio Fernández y M. N. Barrio Rodríguez	79
Indicadores en Patología Digestiva Neutrofilos de la arteria mesentérica superior: una causa infrecuente de obstrucción intestinal J. Sainza-Munizaga, P. Alameda-Serna y J. C. García-Pérez	Presentación intestinal de tuberculosis por endocitocario patológico: diagnóstico asociado a consumo de Bismuto, patología infrecuente y poco conocida Y. Pineda-Vargas, D. M. Acosta y L. A. Alvarez	80
Neumatosis cística intestinal A. F. Romero-Muñoz y R. Barrio-Zelga	Parrot endoscopy: evidence for an achalasia patient with multiple esophageal diverticula Y. Se, H. Zhu and D. Liu	81
Neutrofilos de Weller a pírcula autoinmunitaria C. Ochoa-Sandoval, C. C. Hernández-Segura, J. Pineda-Rodríguez y A. N. González-Fernández	Analisis y endoscopia intestinal: entidades inflamatorias en el diagnóstico diferencial de abdomen agudo C. Sánchez-Jiménez, I. Guevara-Nieto y J. A. Acosta-Paredes	81
Neutrofilos de Weller a pírcula autoinmunitaria C. Ochoa-Sandoval, C. C. Hernández-Segura, J. Pineda-Rodríguez y A. N. González-Fernández	Perforación múltiple de divertículos de intestino delgado en paciente con síndrome de Ulceras Duodenales R. Hernández-Cabrera, A. Burgos-Castro y E. Palencia-López	83
Endoscopic retrieval of trichobezoars in a schizophrenic patient J. L. Bermejo-Hernández, M. E. Torres-Castro and M. Torres-Rodríguez	Altopercia anal: una manifestación paraneoplásica de un adenoma actinómico gástrico J. Barrio Guea, F. Fernández-Serrano y J. de la Fuente-Aguado	83
All that glitters is not gold: A different cause for an "alkaline vomit" A. Pineda, W. Shin, J. Villa-Bueno and G. Navarro	Revisores 2016	85

ARAN

www.reed.es

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

IPD 5477 inglés

Type V biliary cyst with cystolithiasis

Martín Bailón-Cuadrado¹, Pilar Pinto-Fuentes¹, David Pacheco-Sánchez¹ and Trinidad Escudero-Caro²

¹Department of General and Digestive Surgery. Hospital Universitario Río Hortega. Valladolid, Spain. ²Department of Radiology. Hospital Universitario Río Hortega. Valladolid, Spain

Correspondence: Martín Bailón-Cuadrado

e-mail: martin.bc1988@gmail.com

CASE REPORT

We present the case of a 40-year-old female who presented to the General Surgery clinic due to a single episode of abdominal pain which required a visit to the Emergency Department. The patient had undergone surgery during childhood due to the suspicion of a hepatic hydatid cyst. However, an intraoperative cholangiography identified a small, non-complicated biliary cyst. Therefore, a hepatic resection was not performed. The patient did not undergo follow-up of the lesion. A large intrahepatic biliary cyst of 50 x 40 x 35 mm (type V, Todani classification) (1) was identified via ultrasound; it originated from the right hepatic bile duct, with lithiasis inside. This finding was corroborated via nuclear magnetic resonance (NMR) (Figs. 1 and 2). Type V biliary cysts are infrequent and they usually consist of multiple intrahepatic cysts (1). Our patient presented with a large single intrahepatic cyst, which highlights the singularity of this case. In addition, cystolithiasis is a rare finding in biliary cysts, especially in type V (2). Both the hepatic analytical profile and tumor markers were normal. The patient had not suffered previous cholangitis episodes. Despite the fact that type V biliary cysts have a low risk of malignancy, the optimal management of these lesions is surgical resection. Nevertheless, a strict radiologic surveillance was decided upon due to the numerous family related issues that the patient was going

through (3).

Authorship: Every author has participated in the design and development of this article. Every author has participated in the writing and corrections of the manuscript. Every author has approved the final version which will be considered for publication.

Informed consent: The patient authorized the publication of information concerning their medical history as long as it is used for scientific divulgation. Furthermore, we declare that we have followed the official protocols from our medical center to access the contents of medical histories.

REFERENCES

1. Todani T, Watanabe Y, Toki A, et al. Classification of congenital biliary cystic disease: special reference to type Ic and IVA cysts with primary ductal stricture. *J Hepatobiliary Pancreat Surg* 2003;10(5):340-4. DOI: 10.1007/s00534-002-0733-7
2. Söreide K, Körner H, Havnen J, et al. Bile duct cysts in adults. *Br J Surg* 2004;91(12):1538-48. DOI: 10.1002/bjs.4815
3. Sastry AV, Abbadessa B, Wayne MG, et al. What is the incidence of biliary carcinoma in choledochal cysts, when do they develop, and how should it affect management? *World J Surg* 2015;39(2):487-92.




Fig. 1. Coronal view from the NMR showing the large intrahepatic biliary cyst originating from the right hepatic bile duct and placed cranially to gallbladder. A. Intrahepatic biliary cyst originating from the right hepatic bile duct. B. Gallbladder. C. Left hepatic bile duct.




Fig. 2. Reconstruction from the NMR in which the presence of a large amount of lithiasis within the biliary cyst is shown. A. Intrahepatic biliary cyst with multiple lithiasis inside. B. Gallbladder.

Accepted Article